



In This Issue

Potato production updates
Disease forecasting and updates for early and late blight in potato, cucurbit downy mildew updates

Calendar of Events

December 1-3, 2020 – Midwest Food Producers Association Annual Convention/Processing Crops Conference – *Virtual – details to follow*
January 24-26, 2021 – WI Fresh Vegetable Growers Association Educational Conference, Kalahari, Wisconsin Dells, WI (possible remote options)
February 2-4, 2021 – UW-Madison Div. of Extension & WPVGA Grower Education Conference, Holiday Inn, Stevens Point, WI (possible remote options)

Yi Wang, Assistant Professor & Extension Potato and Vegetable Production Specialist, UW-Madison, Dept. of Horticulture, 608-265-4781, Email: wang52@wisc.edu.

This week I am reporting on our 2020 early season russet variety nitrogen trial. We tested Plover Russet and GoldRush in this study, and we had 100, 240, 300, 360, and 420 lb N/acre as the four N treatments. Those total N amount included N from both the fertilizer and the groundwater.




Figure above shows the total yield of GoldRush and Plover Russet under each of the five N rates. Bars with the same letter are not significantly different at $p < 0.05$. For GoldRush, there is no significant difference of total yield between 240, 300, 360 and 420, although the number under 300 lb N/a is the highest, and numbers under 360 and 420 lb N/a are the lowest. For Plover Russet, total yield under 240 lb N/a is significantly higher than those under 360 and 420 lb N/a.



During the growing seasons, we can see clear difference of canopy color under 100 N (left) vs. higher rate at 420 N (right) (photos taken on July 30th).

Figure below shows the yield% of marketable tubers that weighed between 6 oz and 13 oz. 240 lb N/a produced the highest number for GoldRush and 300 lb N/a for Plover Russet.





Figure below shows the yield % of tubers that are lighter than 6 oz. Generally lower N rate at 100 lb N/a produced more smaller tubers.



I took a picture of tubers from the two varieties under each N rate at harvest. We cannot easily eye ball difference of size profiles under different N rates.



Amanda Gevens, Dept. Chair, Professor & Extension Specialist, UW-Madison Plant Pathology, gevens@wisc.edu, Cell: 608-575-3029. <https://vegpath.plantpath.wisc.edu/>

Current P-Day (Early Blight) and Disease Severity Value (Late Blight) Accumulations (Many thanks to Ben Bradford, UW-Madison Entomology; Stephen Jordan, UW-Madison Plant Pathology). A P-Day value of ≥ 300 indicates the threshold for early blight risk and triggers preventative fungicide application. A DSV of ≥ 18 indicates the threshold for late blight risk and triggers preventative fungicide application. Red text in table indicates threshold has been met/surpassed. Weather data used in these calculations comes from weather stations that are placed in potato fields in each of the four locations. Data are available in multiple formats for each station at: <https://vegpath.plantpath.wisc.edu/dsv/>

<i>Location</i>	<i>Planting Date</i>		<i>50% Emergence Date</i>	<i>Disease Severity Values 9/12/2020</i>	<i>Potato Physiological Days 9/12/2020</i>
<i>Grand Marsh</i>	Early	Apr 17	May 18	164	891
	Mid	Apr 25	May 26	161	835
	Late	May 6	June 1	158	795
<i>Hancock</i>	Early	Apr 8	May 18	81	872
	Mid	Apr 20	May 25	79	821
	Late	May 4	May 30	76	783
<i>Plover</i>	Early	Apr 10	May 23	139	826
	Mid	Apr 20	May 30	133	772
	Late	May 5	June 1	133	759
<i>Antigo</i>	Early	May 14	June 5	81	743
	Mid	May 24	June 10	81	705
	Late	Jun 1	June 17	80	658

Late Blight Management: Our DSVs are reported here from emergence to September 12. Over the past week, we saw extremely low accumulations (just 3 DSVs over the course of a week; reminder, max potential DSV per day is 4 under optimum disease conditions). Plantings of potatoes in the Grand Marsh, Hancock, Plover, and Antigo areas should receive routine (~weekly) preventative fungicide application for late blight management. During senescence to harvest, tubers remain susceptible to late blight. In particular, for crops fated for long term storage, continued application of mancozeb-containing fungicide

can provide management of the tuber phase of late blight which can happen during spore movement and 'washout' events. This late season fungicide treatment is especially important if fields are/were proximal to any known infections, and if the maturity of crops varies greatly within your area (due to risk of foliar late blight and spore availability).

Early Blight Management: PDays are exceeding the threshold of 300 for early planted potatoes in Grand Marsh, Hancock, Plover, and Antigo areas. For more information about fungicide selections, please see the Potato section of the A3422 Commercial Vegetable Production Guide for Wisconsin, 2020. <https://cdn.shopify.com/s/files/1/0145/8808/4272/files/A3422-2020.pdf>

National late blight update: No new reports of late blight in Wisconsin. Across the US, there were also no new reports that I am aware of over the past week. On Aug 20 US-23 was confirmed on tomato and potato in Chautauqua County in western NY state. So far this season, there have been just two WI confirmations of late blight from tomato (Pierce Co.) and potato (Adams Co. US-23) this season. No widespread movement from these sites as far as I'm aware. Earlier this season, late blight had been reported in North Carolina (tomato US-23), Washington, and it was found in British Columbia, western Canada (Delta and Surrey) over a month ago now. The site: <https://usablight.org/map/> includes reports as they are submitted in the US. Previous reports documented the disease in AL, GA, NC, NY, FL, TN, WA, and WI. Where the late blight pathogen has been tested in the US so far this year, the clonal lineage has been US-23.

National cucurbit downy mildew update: No downy mildew reported from WI at this time through my Vegetable Pathology Lab or the UW Plant Disease Diagnostic Clinic. Reports to date, have come from: AL, CT, DE, GA, IL, IN, KS, KY, MA, MD, ME, MI, MS, NC, NH, NJ, NY, OH, Ontario & Quebec Canada, PA, SC, TN, VA, and WV. No forecasted movement of the pathogen in our direction, with prevailing air moving eastward. <https://cdm.ipmPIPE.org/forecasting/>